



SILIQ™ REMS Program Patient Wallet Card

SILIQ is indicated for the treatment of moderate to severe plaque psoriasis in adult patients who are candidates for systemic therapy or phototherapy and have failed to respond or have lost response to other systemic therapies.

WARNING: Suicidal thoughts and behavior, including completed suicides, have occurred in patients treated with SILIQ.

Taking SILIQ has proven effective for the treatment of moderate to severe plaque psoriasis in adult patients who are candidates for systemic therapy or phototherapy. However, if you are experiencing sudden feelings of withdrawal, anxiety, depression or hopelessness, call your doctor immediately. Suicide warning signs also include thinking about hurting or killing yourself; seeking access to firearms, pills or other means for the purpose of self-harm; and talking or writing about death and dying when these actions are out of the ordinary.^{1,2}

You are not alone. Help is available.

I will call my doctor or the **National Suicide Prevention Lifeline at 1-800-273-8255 (TALK)** if:

- I feel new or worsening feelings of withdrawal, depression, anxiety, hopelessness, or other mood changes beginning.
- I am thinking about hurting or killing myself; seeking access to firearms, pills or other means for the purpose of self-harm; or am talking or writing about death and dying².

I will **call 911** if I feel an **immediate threat of death or self-injury**.

Learn about the signs of suicide at www.suicidelifeline.org.

For more information, visit www.SILIQREMS.com or call 1-855-511-6135.

¹ American Association of Suicidology. Know the Warning Signs of Suicide. <http://www.suicidology.org/resources/warning-signs>.

² American Foundation for Suicide Prevention. Suicide Warning Signs. <http://www.afsp.org/understanding-suicide/suicide-warning-signs>.